

Supplementary files

The Mechanism of miR-155/miR-15b Axis Contributed to Apoptosis of CD34+ Cells by Upregulation of PD-L1 in Myelodysplastic Syndromes

MeiWan Cao¹, BaoLing Peng², WanFu Xu¹, PeiYu Chen¹, HuiWen Li¹, Yang Cheng¹, Huan Chen¹, LiPing Ye¹, Jing Xie¹, HongLi Wang¹, Lu Ren¹, LiYa Xiong¹, JingNan Zhu³, XiangYe Xu³, LanLan Geng¹ and SiTang Gong¹.

¹ Department of Gastroenterology, Guangzhou Women and Children’s Medical Center, Guangzhou Medical University, Guangzhou City, Guangdong Province, 510623, China.

² Center for child health and mental health, Shenzhen Children’s Hospital, Shenzhen City, Guangdong Province, China.

³ Department of Hematology, Guangzhou Women and Children’s Medical Center, Guangzhou Medical University, Guangzhou City, Guangdong Province, 510623, China.


⁴ Guangzhou Institute of Pediatrics, Guangzhou Women and Children’s Medical Center, Guangzhou Medical University, Guangzhou City, Guangdong Province, 510623, China.

Online Supplement

Supplementary material

Supplementary table 1. Chromosome abnormalities in MDS

Anomaly	total, n (% of all cases)
-5/5q-	6 (15.0)
-7/7q-	4 (10.0)
+8	4 (10.0)
-18/18q-	3 (7.5)
20q-	2 (5.0)
-Y	2 (5.0)
+21	2 (5.0)
-21	1 (2.5)



Supplementary figure. 1 Venetoclax promotes apoptosis of bone marrow CD34+ cells. After treatment of CD34+ cells with the Bcl-2 inhibitor venetoclax, cell proliferation (A), cell cycle (B), apoptosis (C-D), Bcl-2 and Bax protein expression (E) were determined.