

Original Article

Hodgkin Lymphoma in Children: A 16-year Experience at the Children’s Welfare Teaching Hospital of Baghdad, Iraq

Supplementary materials.

Supplemental Table 1. Multivariate COX Regression Overall Survival.

Characteristic	HR ¹	95% CI ¹	p-value
Gender			
Male	—	—	
Female	0.92	0.35, 2.38	0.9
Age			
< 5	—	—	
≥ 5	0.60	0.22, 1.66	0.3
Baghdad			
Baghdad	—	—	
Out	1.93	0.81, 4.57	0.14
Stage class			
IA-IB-IIA	—	—	
IIB-IIIA-IIIB-IVA-IVB	16.0	2.13, 121	0.007
Protocol			
ABVD	—	—	
COPP/ABV/ABVD	1.75	0.70, 4.37	0.2
Other	2.76	0.77, 9.98	0.12

¹HR = Hazard Ratio, CI = Confidence Interval

Supplemental Table 2. Multivariate COX Regression Event-Free Survival.

Characteristic	HR ¹	95% CI ¹	p-value
Gender			
Male	—	—	
Female	0.79	0.47, 1.34	0.4
Age			
< 5	—	—	
≥ 5	0.64	0.35, 1.16	0.14
Baghdad			
Baghdad	—	—	
Out	1.37	0.86, 2.17	0.2
Stage class			
IA-IB-IIA	—	—	
IIB-IIIA-IIIB-IVA-IVB	4.32	2.20, 8.49	<0.001
Protocol			
ABVD	—	—	
COPP/ABV/ABVD	0.53	0.29, 0.98	0.044
Other	1.91	0.94, 3.91	0.075

¹HR = Hazard Ratio, CI = Confidence Interval