



Original article

Integration of CD200, CD43 and ROR1 in Multiparameter Flow Cytometry (MFC) Routine Panels for the Differential Diagnosis of B-cell lymphoproliferative Disorders (B-LPDs)

Supplementary Files

Supplementary Table 1. Patient characteristics, total cohort (n=2615) and according to their B-LPDs diagnosis.

Variables	Total cohort	CLL	MCL	PPBL	CLL/SLL-type MBL	cHCL	vHCL	MZL	FL	DLBCL	Healthy people	LPL	Non-CLL/SLL-type MBL
Total cohort, n (%)	2615	1401 (53.6)	65 (2.5)	16 (0.6)	466 (17.8)	112 (4.2)	32 (1.2)	83 (3.1)	50 (1.9)	34 (1.3)	281 (10.7)	26 (0.9)	32 (1.2)
Age, median years (IQR)	66 (53-74)	69 (59-78)	65 (58-75)	43 (39-52)	57 (48-66)	56.0 (48-68)	69 (62-77)	68 (61-74)	60 (53-68)	72 (62-78)	55 (41-66)	68 (57-76)	72.0 (66-80)
Gender, M/F, n (%)	1509/1106 (57.7/42.3)	810/591 (57.8/42.2)	55/10 (84.6/15.4)	2/14 (12.5/87.5)	255/211 (54.7/45.3)	91/21 (81.2/18.8)	24/8 (74/25)	44/39 (53.0/47.0)	24/26 (48/52)	20/14 (58.8/41.2)	146/135 (51.9/48.1)	17/9 (65.3/34.6)	12/20 (37.5/62.5)
WBC, x10⁹/L, median (IQR)	15.0 (9.98-23.9)	19.5 (15.5-36.2)	14.8 (7.8-34.3)	12.4 (11.0-14.3)	9.8 (8.1-11.4)	3.7 (2.4-6.3)	10.9 (6.5-21.9)	11.0 (7.0-84.5)	16.8 (9.5-34.2)	11.3 (6.4-20.3)	7.6 (5.2-10.5)	10.4 (4.6-17.6)	8.5 (6.2-11.0)
ALC, x10⁹/L, median (IQR)	8.5 (4.6-16.3)	12.8 (9.0-27.4)	9.2 (3.3-15.0)	6.3 (5.3-7.4)	4.7 (3.7-5.0)	2.1 (1.1-3.6)	5.8 (2.7-13.4)	5.9 (3.6-50.1)	10.2 (3.0-20.2)	3.1 (1.2-10.5)	2.7 (1.7-4.2)	4.1 (1.8-6.9)	3.3 (1.9-5.2)
RCL, median % (IQR)	59.7 (48-70.8)	67.8 (59.4-76.2)	55.7 (35.5-66.9)	50.7 (44.3-58.3)	48.9 (42.2-55.0)	57.9 (48.7-66.6)	56.3 (35.6-65.6)	55.5 (42.8-67.9)	56.0 (44.1-75.2)	24.3 (15.1-41.0)	40.8 (30.0-50.1)	42.7 (36.4-59.7)	39.6 (30.9-46.2)
Clonal lymphocyte, x10⁹/L, median (IQR)	6.0 (2.3-13.3)	10.3 (6.5-24.8)	6.8 (2.3-13.2)	-	2.6 (1.5-3.3)	0.5 (0.1-2.1)	5.7 (1.2-12.1)	3.4 (1.8-50.8)	5.9 (1.7-14.9)	2.1 (0.7-7.3)	-	2.1 (0.6-5.0)	0.6 (0.3-1.9)